



converting repulsive d¹⁰-d¹⁰ interactions into partial bonding. With respect to the second question, the design of optimal ligands, an important step was taken by Beck and Strähle recently, who reported a trinuclear Cu(I) complex with the pentaazenido ligand RNNNNNR⁻ (**1**).³




The world's record short Cu(I)-Cu(I) distance, 2.35 Å, is observed in this molecule. In this paper we examine the electronic structure of **1**. We also look at some hypothetical polymers related to this remarkable molecule. But before we do so let us review the structures of a few Cu(I) clusters that also have close Cu-Cu distances.

Gambarotta, Floriani, and co-workers⁴ have recently reported the structure of an interesting pentameric species ([CuMes]₅, **2**).



This structure is shaped like a five-pointed star with the mesityl groups serving as the outer points and the Cu(I) atoms as the inner points of the star. The average Cu(I)-Cu(I) distance is about 2.45 Å, which is 0.1 Å longer than that observed in **1**, but short enough to signal a fairly strong interaction between the Cu(I) atoms.


Earlier Jarvis, Lappert, and co-workers⁵ found the structure of (Me₃SiCH₂Cu)₄ (**3**) to consist of a square planar array of Cu(I)



atoms bridged by alkyl groups. The interesting feature of this compound is the "puckering in" of the Cu(I) atoms, which results in a star-shaped (four points) structure. The distance between the Cu(I) atoms is 2.42 Å, which prior to the discovery of **1** was one of the shortest distances known. Extended Hückel MO calculations done by our group on **3** demonstrated that there is indeed a "soft" bonding interaction between the metal centers.^{2a}


The bonding results from a mixing of the s, p, and d levels, which leads to an attraction between the Cu(I) atoms. Both the stereochemical requirements of the ligand and the direct Cu(I)-Cu(I) interaction contribute to the short metal-metal separation in **3**.

Another structure related to **3** is the one adopted by the mixed transition metal compound 2-cuprio-1-((dimethylamino)methyl)ferrocene (**4**).⁶ This very beautiful structure is formed



by the action of 2-lithio-1-((dimethylamino)methyl)ferrocene on the complex of ((dimethylamino)methyl)ferrocene with copper iodide. Thus, **4** has as its bridging alkyl group a cyclopentadienide dianion moiety, which is then bound in an η⁵ manner to Fe^{II}Cp. The Cu-Cu distance is 2.44 Å.

Several very interesting structures have been observed for the iodocuprates(I) by Hartl and co-workers.⁷ For example, [Cu₅I₇]²⁻ has five CuI₄ tetrahedra condensed in a cyclic array, which leads to the idealized structure **5**.^{7a} However, this idealized structure



would give extraordinarily short Cu(I)-Cu(I) distances (ca. 1.85 Å). The observed structure has the Cu atoms displaced above or below the I₅ plane, yielding a Cu-Cu distance of 2.58 Å, which is about the bond distance observed in copper metal.

Face-shared tetrahedra, an uncommon structural feature in coordination chemistry, are accommodated if the tetrahedra are centered by Cu(I) ions. Two examples, among several, are given by the Cu₂I₃⁻ chain **6**^{8a} and the Cu₃I₄⁻ chain **7**.^{8b} The shortest Cu-Cu separations are 2.45 Å in **6** and 2.46 Å in **7**.


It is clear that in all of these structures there must be some Cu-Cu bonding interaction. The question remains: how much bonding? The 2.35 Å Cu-Cu distance in **1** is spectacularly short, by any measure. The ligand geometry is also highly symmetrical, and so this complex is a natural focus for a further examination of this bond type.

A Simple Picture of Attractive Cu(I) Interactions


Before beginning an analysis of the Beck and Strähle complex **1**, it is useful to review the explanation that we have given for why Cu(I)-Cu(I) interactions are not repulsive.²

- (3) Beck, J.; Strähle, J. *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 409.
 (4) Gambarotta, S.; Floriani, C.; Chiesi-Villa, A.; Guastini, C. *J. Chem. Soc., Chem. Commun.* **1983**, 1156.
 (5) Jarvis, J. A. J.; Kilbourn, B. T.; Pearce, R.; Lappert, M. F. *J. Chem. Soc., Chem. Commun.* **1973**, 475.


- (6) Nesmeyanov, A. N.; Struchkov, Yu. T.; Sedova, N. N.; Andrianov, V. G.; Volgin, Yu. V.; Sazonova, J. *Organomet. Chem.* **1977**, *137*, 217.
 (7) (a) Hartl, H.; Mahdjour-Hassan-Abadi, F. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 378. (b) Mahdjour-Hassan-Abadi, F.; Hartl, H.; Fuchs, J. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 514.
 (8) (a) Andersson, S.; Jagner, S. *Acta. Chem. Scand., Ser. A* **1985**, *A39*, 181. (b) Hartl, H.; Mahdjour-Hassan-Abadi, F. *Z. Naturforsch., B: Inorg. Chem., Org. Chem.* **1984**, *39B*, 149.



If two or more Cu(I) ions were represented well by only their closed d^{10} configurations, one would in fact get only a repulsion between them (8). In one-electron theory this is the consequence




of two-orbital-four-electron repulsive interactions. But on the coppers one also has 4s and 4p valence orbitals. These stabilize both the bonding and antibonding d orbital combinations, as shown schematically in 9.



One can be more specific: The main contributors to Cu–Cu antibonding (if there were only d orbitals on Cu) are the σ and σ^* z^2 combinations shown in 10. Admixture of appropriate symmetry-adapted s and z combinations transforms these, increasing the bonding in σ and decreasing the antibonding in σ^* . In the extreme σ becomes a really good bond and σ^* an innocuous lone-pair combination.

The important feature is an admixture of s and p character into predominantly d orbitals, i.e. hybridization. In fact, in valence bond language one could say that the Cu moves from a d^{10} configuration, which does not allow any bonding, toward a $d^9s^x p^{1-x}$ configuration. Similar things happen in Ni, Pd, and Pt clusters, aggregating eventually to the metal.

Either way, molecular orbital or valence bond, one reaches the essential conclusion that s and p mixing makes for bonding: the




easier such mixing, i.e. the lower down in energy is the s,p relative to d, the more such bonding one will get. So the Cu(I) story is tied to cluster formation in other closed-shell systems, e.g. in group 2 and 12. There it is np mixing into ns that makes for bonding, and one can understand why Be clusters form whereas He ones do not.

Electronic Structure of $\text{Cu}_3(\text{HNNNNNH})_3$

The model we have chosen for 1 is $[\text{Cu}(\text{HNNNNNH})]_3$, with a geometry idealized from the observed structure. The details of the geometry and the extended Hückel calculations used throughout this paper are given in the Appendix. A calculation of this type on the trimer yields a Cu–Cu overlap population of 0.12. So there is a substantial amount of bonding. But where does it come from?

It is natural to think of the Beck and Strähle compound as being formed from an isolated Cu_3^{3+} linear array and three tridentate HNNNNNH⁻ ligands, as in 11. The bonding between the




coppers could be there in the Cu_3^{3+} unit alone, or it could be induced by the ligands. Table I is illuminating in this respect, for it shows the Cu–Cu overlap population, first in the Cu_3^{3+} cluster alone and then in the complex. And in either case we show the effect of including or excluding d or s,p orbitals in the computation.

Looking at the Cu_3^{3+} unit alone, one sees essentially the picture outlined in the preceding section: antibonding between coppers if only d orbitals are included, turning into bonding for the complete valence set. Including the ligands has little effect. The interesting point, though, made by the $\text{Cu}_3(\text{HN}_3\text{H})_3$ “s,p only” entry, is that, even without d orbitals on the copper, the act of binding with the ligand induces some Cu–Cu bonding.

Let us trace down the origins of this intriguing effect and in the process get some insight into the bonding in 1. Because the emphasis is on the s,p bonding we will introduce that first and then the d orbitals. An interaction diagram for the complex 11 is given in Figure 1, but before we discuss it let us approach the components individually.

Cu_3^{3+} is at left. Its orbitals are trivially derived, leading to recognizable “allylic” sets such as those shown in 12. There is







Figure 1. Interaction diagram for 1 (without d orbitals).

Table I. Overlap Populations in Cu_3^{3+} and $\text{Cu}_3(\text{HN}_5\text{H})_3$

cluster	overlap population		
	s,p only	d only	s,p,d
Cu_3^{3+}	(0)	-0.080	+0.132
$\text{Cu}_3(\text{HN}_5\text{H})_3$	+0.055	-0.010	+0.127

also substantial s,p mixing in some of the orbitals. For instance, the low-lying Cu $1a_2''$ orbital is changed into a combination given


schematically by 13. Formally all of these orbitals are empty in Cu_3^{3+} , but we will soon see that they become partially occupied in complexation.



13

For each HNNNNNH^- we expect from a valence formula such as 14 that there are five nitrogen lone pairs and five π orbitals

(three occupied) of pentadienyl type. The π 's are easy to find among the valence orbitals of HN_5H^- (at right in Figure 1). The lone pairs are less obvious, because there is of course some mixing with NN and NH σ bonds, but these also can be located; they are labeled LN_{1-5} in Figure 1.



14


Note that in the free ligand the fifth lone pair combination, LN_5 , is rather high in energy, perilously close to π_4 . One might even predict, if one took the calculation at face value, that the HN_5H^- anion might be a triplet, with an $(\text{LN}_5)^1(\pi_4)^1$ configuration. There is precious little experimental information on the isolated ligand, which is not very stable. Also it could be the calculations are at fault, for the extended Hückel method tends to put σ levels too high relative to π . MNDO calculations⁹ give a larger gap between LN_5 and π_4 . The electronic ambiguity remains interesting, and we will return to some consequences of it when we study some hypothetical polymers. Meanwhile, let us go on to form the complex.

There are three HN_5H^- ligands. They are so far apart in the geometry of **11** that when they are brought together (right side of Figure 1) they interact but little. The threefold symmetry leads to the formation of one nondegenerate and one degenerate orbital from each ligand orbital. Many of the resulting orbitals can form strong interactions with the orbitals from the copper cluster. However, we find, not unexpectedly, that all of the π type ligand orbitals are unable to overlap effectively with any of the copper orbitals. One can see they have approximately or exactly the same energy in the complex as in HN_5H^- .

The ligand lone pairs are stabilized by interaction with empty Cu cluster orbitals. This is, of course, the expected dative bonding. But as a consequence of that mixing, some Cu_3^{3+} cluster orbitals get populated, and those that are Cu–Cu bonding get populated more, because they interact more strongly. They are closer to the HN_5H^- ligand orbitals in energy. The most significant Cu_3^{3+} fragment molecular orbital populations are 0.92 for $1a_1'$ and 0.63 for $1a_2''$. This is how the ligand induces Cu–Cu bonding.

Note that complex formation has "removed" the near degeneracy of LN_5 and π_4 in the free ligand. LN_5 has been stabilized; it becomes the $5e'$ HOMO of the complex. There is about a 0.66-eV gap between HOMO and LUMO in $\text{Cu}_3(\text{HN}_5\text{H})_3$, at this stage in the calculation.

We now return the 3d orbitals to Cu. The changes that happen are small. The d block appears in the region around -14 eV in the calculations (Figure 2 shows the changes in the frontier orbital region). There are some significant interactions that are new, e.g. that shown in **15** between LN_3 and Cu x^2-y^2 . Some π interactions



15

become effective, for instance interchanging the order of the LUMO's $4e''$ and $3a'$. But the major effect of inclusion of Cu 3d as well as 4s and 4p is a small but significant mixing of s, p, and d on the coppers, just as one found in the Cu_3^{3+} case. The Cu_3^{3+} fragment MO's $1a_1'$ and $1a_2''$ are populated by 0.88 and 0.68 electron, respectively, in the complex.

A weak temperature-independent paramagnetism is observed in the complex.³ Beck and Strähle conjectured that might be due to a paramagnetic excited state with a radical anion on the ligand and a Cu^{2+} ion. Our calculated gap is 0.45 eV. This is not big, but our general experience¹⁰ indicates that a gap of that size is





Figure 2. Frontier orbitals for **1** with s,p only (left) and s,p,d (right).

sufficient to guarantee a singlet ground state. Note, incidentally, that promotion of an electron from the $5e'$ HOMO (some Cu–Cu π -bonding character) to the LUMO (ligand π) would not provide a mechanism for strengthening the Cu–Cu bond. We have no explanation for the observed paramagnetism.

Since the Cu–Cu interactions are only weakly attractive, while the Cu–ligand interactions are strong, the main factor determining the record short Cu–Cu separation in **1** is the bite size of the ligand. One is tempted to associate that with the distance between the dentate nitrogens in RN_2R^- . That is about 2.2 Å in the experimental geometry, even shorter than the Cu–Cu spacing. But we must not jump too quickly to an identification on the N...N separation with the true bite size: perhaps it is best to distinguish "geometrical" bite size from an "electronic" one. What matters, we think, is the *overlap* between the orbitals emerging from a ligand and the corresponding metal orbitals.¹¹ The distinction is made clearer, perhaps, in **16**. The geometrical bite size, S_g ,



16

is defined by us as simply the distance between the ligands. But what if the lone pairs don't "point" along a perpendicular to the line between the ligands? The maximum overlap with some metal orbitals might then come at a larger or smaller distance S_e , the electronic bite size. And just to make things complicated, S_e could depend on the metal–ligand separation.

It is not easy to define S_e , since the definition is theoretical and not experimental and one has to specify the metal orbitals interacting with the ligand. The matter merits a separate study, for a variety of ligands. But in the case at hand what we did was to take a single ligand and one Cu atom (s,p,d) placed 2.03 Å

(9) Dewar, M. J. S.; Thiel, W. *J. Am. Chem. Soc.* **1977**, *99*, 4907.

(10) See, e.g., the discussion in: Hay, P. J.; Thibault, J. C.; Hoffmann, R. *J. Am. Chem. Soc.* **1975**, *97*, 4484.

(11) For a review, see: Moore, D. S.; Robinson, S. D. *Adv. Inorg. Chem. Radiochem.* **1986**, *30*, 1.





Figure 3. For a Cu_∞ rod, from left to right: (a) total density of states; (b) z^2 contribution to DOS; (c) xz, yz contribution; (d) s contribution. Parts b-d are magnified by a factor of 3.0.


away from the nitrogen. We then moved it on an arc measured by an angle α (17), keeping Cu-N constant. The optimum α is



1°, pretty close to zero. So in this case the electronic bite size is nearly the same as the geometric one.

Other Oligomers and Polymers

It is obviously possible to imagine four pentaazenido groups around a Cu_3 bar (18). There exist analogous triazenido binuclear complexes $\text{M}_2(\text{RN}_3\text{R})_4$.¹¹ And it is possible to hypothesize extended analogues of 1 and 18 with new ligands, RN_xR , e.g. 19. The problem is not in thinking these molecules up or in doing calculations on them, but in synthesizing them.



Calculations on 18 and 19 yield few surprises. The square-planar crystal field around each Cu is characteristically strong. It pushes one d orbital, $x^2 - y^2$ (z is the Cu...Cu axis) on each center way up in energy. If the electron count were as shown, the destabilized electrons would enter N-N antibonding ligand π^* orbitals. Oxidation of the molecule as a whole (and so of the Cu centers to Cu^{2+} or Cu^{3+}) is more likely.

Polymeric extrapolations of 1 or 18 and 19 spring to mind. The unit cells of these are shown in 20 and 21. The ligand in these






Figure 4. Calculated total DOS and the p_z projection for $(\text{N}_2)_\infty$.

polymers should be formally neutral, if it is to provide the same lone pair and π environment as RN_3R^- and RN_5R^- (22). So the



electron counts one might focus on would be those corresponding to $\text{Cu}(\text{N}_2)_3^+$ in 20, and (given the strong square-planar crystal field) $\text{Cu}(\text{N}_2)_4^{2,3+}$ in 21.

Let us focus first on the Cu_∞ rod that is found in both 20 and 21. The band structure of a linear chain is pretty simple.¹² Decomposition of the density of states (DOS), shown in Figure 3, show the following:

(1) A metal d block, consisting of a wide z^2 or σ band, a narrower xz, yz or π band, and a very undispersed $x^2 - y^2, xy$, or δ band is observed.

(2) There is strong s, p_z mixing, resulting in a Cu-Cu σ bonding band (-8 to -13 eV) and a higher lying σ^* , Cu-Cu antibonding band.¹² The x, y band is much less dispersed, around -6 eV.

(3) There is clear evidence of the same s, p, d mixing that we found so important in Cu_2^{2+} and Cu_3^{3+} : note the "resonances" between z^2 and s contributions at -15 and -13 eV.




Figure 5. Calculated DOS plots for the Cu orbitals, all N orbitals, and $d_{x^2-y^2}$ type orbitals for (a) **20** and (b) **21**. The Fermi level is indicated for unit cell electron counts of 40, 39, and 38 for **20**, while for **21** they are 50, 49, and 48.

Figure 4 shows the computed density of states of the $(N_2)_\infty$ ligand polymer itself, along with the π contributions to it. The wide π band is what one would expect of the polyene analogue.

Tripling or quadrupling this polymer, as one must to form **20** or **21**, will not do much to the DOS. But interaction with the Cu chain has more serious effects. Selected contributions to the DOS for the composite polymers **20** and **21** are shown in Figure 5.

The following points may be noted:

(1) These molecules should be conducting, for most reasonable electron counts. The possibility exists, however, that for electron counts close to Cu^{2+} , these polymers might be antiferromagnetically coupled, as in the familiar copper acetates.¹³

(2) The Fermi level comes in a region of mainly $(N_2)_\infty$ ligand states, both σ (lone pair) and π . Recall that even in the $Cu_3-(RN_3R)_3$ complex there was already a small gap between π_4 and $LN_5 + \lambda Cu$ combinations. As the ligand chain is extended that gap must vanish—the π levels and the lone pair levels develop into a band.

(3) Crystal field effects are apparent: note the low energy of the $x^2 - y^2$ contribution in the trigonally coordinated $Cu(N_2)_3$ polymer, contrasted with the high energy of these orbitals in the

square-planar $Cu(N_2)_4$ environment.

Acknowledgment. We thank J. Strähle for the impetus for this work, and for providing some experimental information prior to publication. K.M.M. would like to thank Ralph Wheeler, Marja Zonneville, and Jing Li for enlightening conversations regarding this research. Jing Li is to be especially thanked for reproducing some of the calculations at Cornell and sending the results over BITNET to Lund University, Lund, Sweden. We are grateful to the Swedish National Research Council for a Tage Erlander Professorship, which made our stay at the Kemcentrum in Lund possible. We thank Prof. Sten Andersson and his group for their hospitality. Our research at Cornell was generously supported by the National Science Foundation through Research Grant DMR821722702 to the Materials Science Center. We are grateful to Jane Jorgensen and Elisabeth Fields for their expert drawings.

Appendix

The extended Hückel method¹⁴ in the tight binding approximation¹⁵ was used in all calculations. The parameters used for carbon, hydrogen, and copper were as follows: $H_{ss} = -26.0$ eV, $H_{pp} = -13.4$ eV, $\zeta_{s,p} = 1.95$; $H_{ss} = -13.6$ eV, $\zeta_{s,p} = 1.3$; $H_{ss} = -11.4$

(13) For a leading reference, see: Harcourt, R. D.; Skrezenek, F. L.; Macclagan, R. G. A. *R. J. Am. Chem. Soc.* **1986**, *108*, 5403.

(14) Hoffmann, R. *J. Chem. Phys.* **1963**, *39*, 1397. Hoffmann, R.; Lipscomb, W. N. *J. Chem. Phys.* **1963**, *36*, 3179; **1962**, *37*, 2872.

(15) Whangbo, M.-H.; Hoffmann, R. *J. Am. Chem. Soc.* **1978**, *100*, 6093.

eV, $H_{pp} = -6.06$ eV, $H_{dd} = -14.0$ eV, $\zeta_{s,p} = 2.2$, $\zeta_{d1} = 5.95$, $\zeta_{d2} = 2.3$, $c_1 = 0.5933$, $c_2 = 0.5744$ respectively. To calculate average properties for the 1-D cases a 100-point k-point set was used.¹⁶ The geometries used for calculations were as follows: Cu-Cu-

(monomer), 2.35 Å; Cu-Cu(polymer), 2.26 Å; N-N, 1.34 Å; Cu-N, 2.03 Å; N-H, 1.0 Å; Cu-N-N, 122.5°; N-N-N, 115.0°; H-N-N, 118.0°.

Registry No. [Cu(tolylNNNNNtolyl)]₃, 96129-16-7; [Cu₃(tolylNNNNNtolyl)₄]⁻, 114326-40-8; [Cu(tolylNNNNNtolyl)]₄, 114350-60-6.

(16) Pack, J. D.; Monkhorst, J. H. *Phys. Rev. B: Solid State* 1977, 16, 1748.

Contribution from the Department of Chemistry and Laboratory for Molecular Structure and Bonding, Texas A&M University, College Station, Texas 77843

Preparation and Properties of the Mo₂X₄(PMe₃)₄ Compounds with X = NCS and NCO

F. Albert Cotton* and Marek Matusz

Received January 20, 1988

The Mo₂X₄(PMe₃)₄ compounds where X = NCO and NCS are formed in a high-yield, one-pot reaction from Mo₂(O₂CCH₃)₄ and (CH₃)₃SiX in the presence of the phosphine. Both compounds crystallize in the tetragonal space group *P*4₂1₂ with the following unit cell dimensions. X = NCO: $a = b = 11.990$ (2) Å, $c = 20.271$ (2) Å, $V = 2914$ (1) Å³, $Z = 4$. X = NCS: $a = b = 12.556$ (1) Å, $c = 21.033$ (4) Å, $V = 3316$ (1) Å³, $Z = 4$. Both molecules consist of quadruply bonded molybdenum atoms with trans phosphines and X groups as ligands. Electrochemical and spectroscopic properties of these compounds have been investigated and correlated with other compounds where X = Cl, Br, and I.

Introduction

Mo₂X₄L₄ complexes are among the most often studied multiple-bonded dimetal complexes. They have contributed much to the understanding of the nature of the multiple bonds between metal atoms.¹ In the past, conclusions have been drawn on some incomplete classes of these compounds. Recently the discovery of some new compounds, namely Mo₂I₄(PMe₃)₄² and Mo₂I₄(dppe)₂,³ has shed some new light on the subject. The Mo₂X₄(PMe₃)₄ (X = Cl, Br, I) set has received the most attention, because of the simplicity of the system, but two other classes, Mo₂X₄(dppe)₂ (X = Cl, Br, I) and Mo₂X₄(dppm)₂ (X = Cl, Br, I, NCS), have also been investigated and structurally characterized. In view of their theoretical importance, we decided to extend the Mo₂X₄(PMe₃)₄ series with some new compounds. As a part of our recent interest⁴ in the thiocyanate chemistry of M₂ species, we have prepared Mo₂(NCS)₄(PMe₃)₄. By the same synthetic route Mo₂(NCO)₄(PMe₃)₄ was also prepared. These two compounds, together with the Cl, Br, and I derivatives, form the largest known class of quadruply bonded, homologous compounds. In this study we report the preparation and molecular structures of these two new compounds, as well as their spectroscopic and electrochemical properties. The relationship between the energy of the δ - δ^* transition and the first reduction potential for the entire class has been examined.

Experimental Section

All reactions were done under anaerobic conditions. Me₃SiNCO and Me₃SiNCS were purchased from Aldrich Chemical Co. Trimethylphosphine was purchased from Strem Chemicals Inc.

Preparation of Mo₂(NCO)₄(PMe₃)₄ (1). A round-bottom flask was charged with 1.0 g (2.3 mmol) of Mo₂(O₂CCH₃)₄, 0.858 g (11 mmol) of PMe₃, 1.26 g (11 mmol) of Me₃SiNCO, and 50 mL of toluene. The reaction mixture was stirred at room temperature for 2 days and then evaporated to dryness. The solid residue was dissolved in hot toluene (about 150 mL), and the mixture was filtered hot through a short Celite column. The solution was refrigerated overnight to give well-formed crystals of the product, which were filtered out, washed with hexane, and vacuum-dried. The yield was 1.2 g (72% based on Mo) of blue-black

crystals. They are soluble in THF, benzene, and CH₂Cl₂ to give sky blue solutions. The compound can be handled in the air in the solid state but decomposes slowly in solution. UV-vis (CH₂Cl₂ solution): $\lambda = 617$ nm ($\epsilon = 8050$ M⁻¹ cm⁻¹), $\lambda = 460$ nm ($\epsilon = 430$ M⁻¹ cm⁻¹), $\lambda = 330$ nm ($\epsilon = 12370$ M⁻¹ cm⁻¹). The same λ_{max} values were recorded for the THF solution of the compound. IR (cm⁻¹): 2210 vs, 1380 m, 1345 m, 1305 m, 1290 m, 960 s, 865 w, 855 w, 750 m, 680 m, 620 m, 380 m. ³¹P NMR: -8.30 ppm (s).

Mo₂(NCS)₄(PMe₃)₄ (2). The experimental procedure is essentially the same as that described above. Isolated yields were typically 80% of a dark green crystalline solid. Solubility and stability are similar to those of the NCO derivative. UV-vis (CH₂Cl₂ solution): $\lambda = 670$ nm ($\epsilon = 6680$ M⁻¹ cm⁻¹), $\lambda = 497$ nm ($\epsilon = 380$ M⁻¹ cm⁻¹), $\lambda = 395$ nm ($\epsilon = 23100$ M⁻¹ cm⁻¹). IR (cm⁻¹): 2040 vs, 1380 m, 1310 m, 1290 m, 960 s, 870 m, 810 m, 805 m, 750 m, 745 m, 625 m, 490 m, 360 m. ³¹P NMR: -6.8 ppm (s). Anal. Calcd for Mo₂S₄P₄N₄C₁₆H₃₆: C, 26.40; H, 4.98; N, 7.69. Found: C, 26.65; H, 5.08; N, 7.54.

Mo₂(NCS)₄(PMe₃)₄ could also be prepared by an alternative procedure: Mo₂Cl₄(PMe₃)₄, 0.20 g (0.313 mmol), KSCN, 0.122 g (1.25 mmol), and 15 mL of MeOH were refluxed for 24 h. The reaction mixture turned from fluorescent red-blue to green during this time. It was cooled and filtered. The solid residue was washed with a small amount of MeOH and extracted with boiling toluene. The toluene solution was filtered hot through a Celite bed and refrigerated. Next day the crystalline product was filtered out and vacuum-dried. The yield was 0.144 g (53%). The spectroscopic (UV-vis, ³¹P NMR) and electrochemical properties were identical with those of the sample obtained by the Me₃SiNCS method.

We were unable to obtain Mo₂(NCO)₄(PMe₃)₄ by the similar ligand-exchange reaction.

Measurements. Elemental analyses were performed by Galbraith Laboratories Inc. IR spectra were recorded as Nujol mulls on KBr plates (4000-600 cm⁻¹) or CsI plates (600-300 cm⁻¹) with a Perkin-Elmer 785 spectrophotometer. The electronic spectra were recorded on dichloromethane solutions (Aldrich, HPLC grade) with a Cary 17D spectrophotometer. Electrochemical measurements were done with a BAS-100 electrochemical analyzer, employing 0.2 M tetra-*n*-butylammonium hexafluorophosphate solutions in THF or CH₂Cl₂. A three-electrode configuration was used with platinum working and auxiliary electrodes and a silver-silver chloride reference electrode. The potentials are corrected for the solution resistance. Under the experimental conditions, ferrocene is oxidized at $E_{1/2} = 0.510$ V in CH₂Cl₂ and $E_{1/2} = 0.595$ V in THF. ³¹P NMR samples were recorded on CDCl₃ solutions with a Varian XL 200 spectrometer and are referenced to 85% H₃PO₄.

X-ray Crystallography

Single crystals of compounds 1 and 2, obtained as described in the experimental section, were glued on top of glass fibers. Indexing revealed tetragonal cells, Laue class (*4/mmm*), and axial dimensions were con-

- (1) Cotton, F. A.; Walton, R. A. *Multiple Bonds Between Metal Atoms*; Wiley: New York, 1982.
- (2) Hopkins, M. D.; Schaefer, W. D.; Bronikowski, M. J.; Woodruff, W. H.; Miskowski, V. M.; Dallinger, R. F.; Gray, H. B. *J. Am. Chem. Soc.* 1987, 109, 408. Cotton, F. A.; Poli, R. *Inorg. Chem.* 1987, 26, 3228.
- (3) Cotton, F. A.; Dunbar, K. R.; Matusz, M. *Inorg. Chem.* 1986, 25, 3641.
- (4) Cotton, F. A.; Matusz, M. *Inorg. Chem.* 1987, 26, 3468.